

Some Quantitative Characterizations of Certain Symplectic Groups

M. Akbari and A. R. Moghaddamfar

*Department of Mathematics, K. N. Toosi University of Technology,
P. O. Box 16315-1618, Tehran, Iran*

E-mails: moghadam@kntu.ac.ir and moghadam@ipm.ir

April 30, 2013

Abstract

Given a finite group G , denote by $D(G)$ the degree pattern of G and by $OC(G)$ the set of all order components of G . Denote by $h_{OD}(G)$ (resp. $h_{OC}(G)$) the number of isomorphism classes of finite groups H satisfying conditions $|H| = |G|$ and $D(H) = D(G)$ (resp. $OC(H) = OC(G)$). A finite group G is called OD-characterizable (resp. OC-characterizable) if $h_{OD}(G) = 1$ (resp. $h_{OC}(G) = 1$). Let $C = C_p(2)$ be a symplectic group over binary field, for which $2^p - 1 > 7$ is a Mersenne prime. The aim of this article is to prove that $h_{OD}(C) = 1 = h_{OC}(C)$.

Keywords: spectrum of a group, prime graph, degree pattern, order component, symplectic group $C_n(q)$, OD(OC)-characterizability of a finite group.

1 Introduction

Only finite groups will be considered. Let G be a group, $\pi(G)$ the set of all prime divisors of its order and $\omega(G)$ be the spectrum of G , that is the set of its element orders. The *prime graph* $GK(G)$ (or *Gruenberg-Kegel graph*) of G is a simple graph whose vertex set is $\pi(G)$ and two distinct vertices p and q are joined by an edge (and we write $p \sim q$) if and only if $pq \in \omega(G)$. Let $t(G)$ be the number of connected components of $GK(G)$. The i th connected component is denoted by $\pi_i(G)$ for each $i = 1, 2, \dots, t(G)$. In the case when $2 \in \pi(G)$, we assume that $2 \in \pi_1(G)$.

The classification of simple groups with disconnected prime graph was obtained by Williams [26] and Kondratév [12]. Moreover, a corrected list of these groups can be found in [13]. Recall that a *clique* in a graph is a set of pairwise adjacent vertices. Note that for all non-abelian simple groups S with disconnected prime graph, all connected components $\pi_i(S)$ for $2 \leq i \leq t(S)$ are clique, for instance, see [12], [23] and [26].

The *degree* $\deg_G(p)$ of a vertex $p \in \pi(G)$ in $GK(G)$ is the number of edges incident on p . If $\pi(G) = \{p_1, p_2, \dots, p_h\}$ with $p_1 < p_2 < \dots < p_h$, then we define

$$D(G) := (\deg_G(p_1), \deg_G(p_2), \dots, \deg_G(p_h)),$$

which is called the *degree pattern* of G .

Given a group G , denote by $h_{OD}(G)$ the number of isomorphism classes of groups with the same order and degree pattern as G . For instance, if G is a cyclic group of order p , where p is a prime number, then $h_{OD}(G) = 1$. Notice that the cyclic group $\mathbb{Z}_p$ is the only group of order p . Similarly, in the case when G is a p -group of order p^2 , $h_{OD}(G) = 2$, in fact, the non-isomorphic groups $\mathbb{Z}_{p^2}$ and

$\mathbb{Z}_p \times \mathbb{Z}_p$ are the only groups with the same order and degree pattern. All finite groups, in terms of the function $h_{OD}(\cdot)$, are classified as follows:

Definition 1 A group G is called k -fold OD-characterizable if $h_{OD}(G) = k$. Usually, a 1-fold OD-characterizable group is simply called OD-characterizable.

There are scattered results in the literature showing that certain simple groups are k -fold OD-characterizable for $k \in \{1, 2\}$ (see Table 1). In this article, we will focus our attention on the symplectic groups $C_p(2) \cong S_{2p}(2)$, where p is an odd prime. Recall that $C_2(2)$ is not a simple group, in fact, the derived subgroup $C_2(2)'$ is a simple group which is isomorphic with $A_6 \cong L_2(9)$. In addition, we recall that $B_2(3) \cong {}^2A_4(2^2)$, $B_n(2^m) \cong C_n(2^m)$ and $B_2(q) \cong C_2(q)$ (see [4]). Previously, it was determined the values of $h_{OD}(\cdot)$ for some symplectic and orthogonal groups (see [2, 14]):

G	Restrictions on G	$h_{OD}(G)$
$B_3(4) \cong C_3(4)$		1
$B_2(q) \cong C_2(q)$	$ \pi(\frac{q^2+1}{2, q-1}) = 1$	1
$B_{2^m}(q) \cong C_{2^m}(q)$	$ \pi(\frac{q^2+1}{2, q-1}) = 1$, q is even	1
$B_3(5), C_3(5),$		2
$B_n(q), C_n(q),$	$n = 2^m > 2$, $ \pi(\frac{q^n+1}{2}) = 1$, q is an odd prime power	2
$B_p(3), C_p(3),$	$ \pi(\frac{3^p-1}{2}) = 1$, p is an odd prime,	2

If n is a natural number, then $\pi(n)$ denotes the set of all prime divisors of n . Given a group G , the order of G can be expressed as a product of some co-prime natural numbers $m_i(G)$, $i = 1, 2, \dots, t(G)$, with $\pi(m_i(G)) = \pi_i(G)$. The numbers $m_1(G), m_2(G), \dots, m_{t(G)}(G)$ are called the *order components* of G . We set

$$OC(G) := \{m_1(G), m_2(G), \dots, m_{t(G)}(G)\}.$$

A list of order components of simple groups with disconnected prime graphs can be found in [5, Tables 1-4]. In the similar manner, we define $h_{OC}(G)$ as the number of isomorphism classes of finite groups with the same set $OC(G)$ of order components. Again, in terms of function $h_{OC}(\cdot)$, the groups G are classified as follows:

Definition 2 A finite group G is called k -fold OC-characterizable if $h_{OC}(G) = k$. In the case when $k = 1$ the group G is simply called OC-characterizable.

It is worth mentioning that the characterization of finite groups through their order components was first introduced by Chen in [5]. A Mersenne prime is a prime that can be written as $2^p - 1$ for some prime p . The purpose of this article is to prove the following theorem.

Theorem 1 Let $C = C_p(2)$ be the symplectic group over binary field, for which $2^p - 1 > 7$ is a Mersenne prime. Then $h_{OD}(C) = 1 = h_{OC}(C)$.

Remark. It is worth noting that the values of functions $h_{OD}(\cdot)$ and $h_{OC}(\cdot)$ may be different. For instance, suppose $M \in \{B_3(5), C_3(5)\}$. By [26], the prime graph associated with M is connected and so $OC(M) = \{|M| = 2^9 \cdot 3^4 \cdot 5^9 \cdot 7 \cdot 13 \cdot 31\}$. On the other hand, it is easy to see that the prime graph associated with a nilpotent group is always a clique, hence, we have

$$h_{OC}(M) > \nu_{\text{nil}}(|M|) \geq \nu_a(|M|) = \text{Par}(9)^2 \cdot \text{Par}(4) = 30^2 \times 5 = 4500,$$

where $\nu_{\text{nil}}(n)$ (resp. $\nu_a(n)$) signifies the number of non-isomorphic nilpotent (resp. abelian) groups of order n and $\text{Par}(n)$ denotes the number of partitions of n . However, by Theorem 1.3 in [2], we know that $h_{OD}(M) = 2$.

We now introduce some further notation. If a is a natural number, r is an odd prime and $(r, a) = 1$, then by $e(r, a)$ we denote the multiplicative order of a modulo r , that is the minimal natural number n with $a^n \equiv 1 \pmod{r}$. If a is odd, we put

$$e(2, a) = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{4}, \\ 2 & \text{if } a \equiv -1 \pmod{4}. \end{cases}$$

We also define the function $\eta : \mathbb{N} \rightarrow \mathbb{N}$, as follows

$$\eta(m) = \begin{cases} m & \text{if } m \equiv 1 \pmod{2}, \\ \frac{m}{2} & \text{if } m \equiv 0 \pmod{2}. \end{cases}$$

Moreover, we will use the notation $\mathbb{A}_n$ and $\mathbb{S}_n$ to denote the alternating and the symmetric group on n letters, respectively. All unexplained notation and terminology are borrowed from the Atlas [7].

2 Preliminaries

The following lemma is a consequence of Zsigmondy's theorem (see [37]).

Lemma 1 *Let a be a natural number greater than 1. Then for every natural number n there exists a prime r with $e(r, a) = n$ but for the cases $(n, a) \in \{(1, 2), (1, 3), (6, 2)\}$*

A prime r with $e(r, a) = n$ is called a *primitive prime divisor* of $a^n - 1$. By Lemma 1, such a prime exists except for the cases mentioned in the lemma. Given a natural number a , we denote by $\text{ppd}(a^n - 1)$ the set of all primitive prime divisors of $a^n - 1$. By our definition, we have $\pi(a - 1) = \text{ppd}(a - 1)$ but for the following sole exception, namely, $2 \notin \text{ppd}(a - 1)$ if $e(2, a) = 2$. In this case, we assume that $2 \in \text{ppd}(a^2 - 1)$.

Lemma 2 ([25]) *Let M be one of the simple groups of Lie type, $B_n(q)$ or $C_n(q)$, over a field of characteristic p . Let r, s be odd primes with $r, s \in \pi(M) \setminus \{p\}$. Suppose that $r \in \text{ppd}(q^k - 1)$, $s \in \text{ppd}(q^l - 1)$ and $1 \leq \eta(k) \leq \eta(l)$. Then r and s are non-adjacent if and only if $\eta(k) + \eta(l) > n$ and $\frac{l}{k}$ is not an odd natural number.*

Lemma 3 ([24]) *Let M be one of the simple groups of Lie type, $B_n(q)$ or $C_n(q)$, over a field of characteristic p , and let $r \in \pi(M) \setminus \{p\}$ and $r \in \text{ppd}(q^k - 1)$. Then r and p are non-adjacent if and only if $\eta(k) > n - 1$.*

Using Lemmas 2 and 3, we have:

- The prime graphs $\text{GK}(B_n(q))$ and $\text{GK}(C_n(q))$ coincide [24, Proposition 7.5].
- $|B_n(q)| = |C_n(q)|$ and $D(B_n(q)) = D(C_n(q))$.

Corollary 1 *Let $p > 3$ be a prime and $C = C_p(2)$. Then $\deg_C(3) = |\pi_1(C)| - 1$.*

Proof. Recall that

$$\pi_1(M) = \pi\left(2(2^p + 1) \prod_{i=1}^{p-1} (2^{2^i} - 1)\right) \quad \text{and} \quad \pi_2(M) = \pi(2^p - 1).$$

Moreover, by Lemma 2, it follows that only primitive prime divisors of $2^p - 1$ are non-adjacent to 3. But by Lemmas 2 and 3 we deduce that $\deg(3) = |\pi_1(M)| - 1$, as desired. $\square$

The following easy lemma (which is appeared in [6]) is crucial to the study of characterizability of symplectic groups $C_p(2)$ by order components.

Lemma 4 *Let G be a group whose prime graph has more than one component. If H is a normal π_k -subgroup of G , then $|H| - 1$ is divisible by m_i , $i \neq k$.*

Lemma 5 ([15]) *Let S be a simple group with a disconnected prime graph $\text{GK}(S)$, except $U_4(2)$ and $U_5(2)$. If G is a group with $\text{OC}(G) = \text{OC}(S)$, then G is neither a Frobenius group nor a 2-Frobenius group.*

Lemma 6 ([26]) *Let G be a group with $t(G) \geq 2$. Then one of the following hold:*

- (1) G is either a Frobenius group or a 2-Frobenius group.
- (2) G has a normal series $1 \trianglelefteq H \triangleleft K \trianglelefteq G$ such that H is a nilpotent π_1 -group, K/H is a non-abelian simple group, G/K is a π_1 -group, $|G/K|$ divides $|\text{Out}(K/H)|$ and any odd order component of G is equal to one of the odd order components of K/H .

Lemma 7 ([8]) *The only solution of the equation $p^m - q^n = 1$, where p, q are primes and $m, n > 1$ are integers, is $(p, q, m, n) = (3, 2, 2, 3)$.*

Given a natural number B and a prime number t , we denote by B_t the t -part of B , that is the largest power of t dividing B .

Lemma 8 ([22]) *Let $B = (2^2 - 1)(2^4 - 1) \cdots (2^{2^n} - 1)$. If t is a prime divisor of B , then $B_t < 2^{3n}$. Furthermore, if $t \geq 5$ then $B_t < 2^{2n}$.*

3 Proof of Theorem 1

Throughout this section, we will assume that $2^p - 1 > 7$ is a Mersenne prime and $C = C_p(2)$. Suppose that G is a group with the same order and degree pattern as C , that is

$$|G| = |C| = 2^{p^2} \prod_{i=1}^p (2^{2^i} - 1) \quad \text{and} \quad D(G) = D(C).$$

Note that, according to the results summarized in [12], we have

$$\pi_1(C) = \pi \left(2(2^p + 1) \prod_{i=1}^{p-1} (2^{2^i} - 1) \right) \quad \text{and} \quad \pi_2(C) = \{2^p - 1\}.$$

By our hypothesis, it is easy to see that

$$\pi_2(G) = \pi_2(C) = \{2^p - 1\} \quad \text{and} \quad \pi(G) = \pi(C) = \pi_1(C) \cup \{2^p - 1\},$$

and so $t(C) = 2$. First of all, we notice that $2^p - 1$ is the largest prime in $\pi(G) = \pi(C)$. Moreover, it follows from Corollary 1 that

$$\deg_G(3) = \deg_C(3) = |\pi_1(C)| - 1,$$

and this forces $\pi_1(G) = \pi_1(C)$ and $t(G) = 2$. Hence, we have

$$\text{OC}(G) = \text{OC}(C) = \left\{ 2^{p^2} (2^p + 1) \prod_{i=1}^{p-1} (2^{2^i} - 1), \quad 2^p - 1 \right\},$$

and from Lemma 5, the group G is neither a Frobenius group nor a 2-Frobenius group. Finally, Lemma 6, reduces the problem to the study of the simple groups. Indeed, by Lemma 6, there is a normal series $1 \trianglelefteq H \triangleleft K \trianglelefteq G$ of G such that:

- (1) H is a nilpotent π_1 -group, K/H is a non-abelian simple group and G/K is a π_1 -group. Moreover, we have

$$K/H \leq G/H \leq \text{Aut}(K/H),$$

and $t(K/H) \geq t(G) \geq 2$,

- (2) $2^p - 1$ is the only odd order component of G which is equal to one of those of the quotient K/H ,
(3) $|G/K|$ divides $|\text{Out}(K/H)|$.

Now we will continue the proof step by step.

Step 1 $K/H \not\cong {}^2A_3(2), {}^2F_4(2)', {}^2A_5(2), E_7(2), E_7(3), A_2(4), {}^2E_6(2)$ or one of the sporadic simple groups.

Note that either the odd order components of above groups are not equal to a Mersenne prime $2^p - 1 > 7$ or their orders does not divide the order of G .

Step 2 $K/H \not\cong \mathbb{A}_n$, where n and $n - 2$ are both prime numbers.

In this case, it follows that $n = 2^p - 1$. Now, simple computations show that

$$\left(\frac{n!}{2}\right)_2 = 2^{\left(\left[\frac{n}{2}\right] + \left[\frac{n}{2^2}\right] + \dots\right) - 1} = 2^{2^p - p - 2}.$$

Now, if $p > 5$, then $2^p - p - 2 > p^2$ and hence the 2-part of $|A_n|$ does not divide the 2-part of $|G|$, a contradiction. In the case when $p = 5$, then $n = 31$ and $|K/H| = (31!)/2$, which does not divide $|G| = |C_5(2)| = 2^{25} \cdot 3^6 \cdot 5^2 \cdot 7 \cdot 11 \cdot 17 \cdot 31$, again a contradiction.

Step 3 $K/H \not\cong \mathbb{A}_n$, where $n = q, q + 1, q + 2$ (q is a prime number), and one of $n, n - 2$ is not prime.

Here, $\{q\}$ is the only odd order component of K/H , and so $q = 2^p - 1$. Now, we consider the alternating group $\mathbb{A}_q$ which is a subgroup of $K/H \cong \mathbb{A}_n$. Similar arguments as those in the previous step, on the subgroup $\mathbb{A}_q$ instead of $\mathbb{A}_n$, lead us a contradiction.

Step 4 K/H is isomorphic neither ${}^2E_6(q)$, $q > 2$, or $E_6(q)$.

We deal with ${}^2E_6(q)$, $q > 2$, the proof for $E_6(q)$ being quite similar. Suppose $K/H \cong {}^2E_6(q)$. First of all, we recall that

$$|{}^2E_6(q)| = \frac{1}{(3, q + 1)} q^{36} (q^{12} - 1)(q^9 + 1)(q^8 - 1)(q^6 - 1)(q^5 + 1)(q^2 - 1).$$

Considering the only odd order component of ${}^2E_6(q)$, that is $(q^6 - q^3 + 1)/(3, q + 1)$, we must have $(q^6 - q^3 + 1)/(3, q + 1) = 2^p - 1$, which implies that $q^9 > 2^p$, or equivalently $q^{36} > 2^{4p}$. Let $q = r^f$. If r is an odd prime, then from Lemma 8, we get

$$q^{36} = r^{36f} = |K/H|_r \leq |G|_r < 2^{3p},$$

which is a contradiction. Therefore we may assume that $r = 2$. In this case, we have $(2^{6f} - 2^{3f} + 1)/(3, 2^f + 1) = 2^p - 1$. Now, if $(3, 2^f + 1) = 1$, then we obtain $2^{3f}(2^{3f} - 1) = 2(2^{p-1} - 1)$, from which we deduce that $3f = 1$, a contradiction. In the case where $(3, 2^f + 1) = 3$, an easy calculation shows that $2^{3f}(2^{3f} - 1) = 2^2(3 \cdot 2^{p-2} - 1)$, and so $3f = 2$, which is again a contradiction.

Step 5 $K/H \not\cong F_4(q)$, where q is an odd prime power.

We remark that $q^4 - q^2 + 1$ is the only odd order component of $F_4(q)$, and clearly this forces $q^4 - q^2 + 1 = 2^p - 1$. Then $q^2(q^2 - 1) = 2(2^{p-1} - 1)$, which shows that $2(2^{p-1} - 1)$ is divisible by 4, a contradiction.

Step 6 $K/H \not\cong {}^2B_2(q)$, where $q = 2^{2m+1} > 2$.

Recall that $|{}^2B_2(q)| = q^2(q^2 + 1)(q - 1)$ and the odd order components of ${}^2B_2(q)$ are

$$q - 1, \quad q - \sqrt{2q} + 1, \quad q + \sqrt{2q} + 1.$$

If $q - 1 = 2^p - 1$, then $q = 2^p$. Now, we consider the primitive prime divisor $r \in \text{ppd}(2^{4p} - 1)$. Clearly $r \in \pi(2^{2p} + 1)$, and so $r \in \pi({}^2B_2(q)) \subseteq \pi(G)$. This is a contradiction.

In the case when $q - \sqrt{2q} + 1 = 2^p - 1$ (resp. $q + \sqrt{2q} + 1 = 2^p - 1$), by simple computations we obtain $2^{m+1}(2^m - 1) = 2(2^{p-1} - 1)$ (resp. $2^{m+1}(2^m + 1) = 2(2^{p-1} - 1)$), a contradiction.

Step 7 $K/H \not\cong E_8(q)$, where $q \equiv 2, 3 \pmod{5}$.

The odd order components of $E_8(q)$ in this case are

$$q^8 - q^4 + 1, \quad \frac{q^{10} + q^5 + 1}{q^2 + q + 1}, \quad \frac{q^{10} - q^5 + 1}{q^2 - q + 1}.$$

If $q^8 - q^4 + 1 = 2^p - 1$, then we obtain $q^4(q - 1)(q + 1)(q^2 + 1) = 2(2^{p-1} - 1)$. However, the left hand side is divisible by 16, while the right hand side is divisible by 2, an impossible.

If $(q^{10} + q^5 + 1)/(q^2 + q + 1) = 2^p - 1$, then after subtracting 1 from both sides of this equation and some simple computations, we obtain

$$q(q - 1)(q + 1)(q^2 + 1)(q^3 + q^2 - 1) = 2(2^{p-1} - 1).$$

Now, if q is odd, then the left hand side is divisible by 16, a contradiction. Moreover, if q is even, then it follows that $q = 2$, and if this is substituted in above equation we get $83 = 2^{p-2}$, a contradiction.

The case $(q^{10} - q^5 + 1)/(q^2 - q + 1) = 2^p - 1$ is quite similar to the previous case and it is omitted.

Step 8 $K/H \not\cong E_8(q)$, where $q \equiv 0, 1, 4 \pmod{5}$.

The odd order components of $E_8(q)$ in this case are

$$\frac{q^{10} + 1}{q^2 + 1}, \quad q^8 - q^4 + 1, \quad \frac{q^{10} + q^5 + 1}{q^2 + q + 1}, \quad \frac{q^{10} - q^5 + 1}{q^2 - q + 1}.$$

Consider the first case. Let $(q^{10} + 1)/(q^2 + 1) = 2^p - 1$. Subtracting 1 from both sides of this equality, we get

$$q^2(q^2 - 1)(q^4 + 1) = 2(2^{p-1} - 1),$$

which implies $2(2^{p-1} - 1)$ is divisible by 4, a contradiction.

Similarly, if $q^8 - q^4 + 1 = 2^p - 1$, we obtain $q^4(q - 1)(q + 1)(q^2 + 1) = 2(2^{p-1} - 1)$, which shows that $2(2^{p-1} - 1)$ is divisible by 16, a contradiction.

Similar arguments work if $(q^{10} + q^5 + 1)/(q^2 + q + 1) = 2^p - 1$ or $(q^{10} - q^5 + 1)/(q^2 - q + 1) = 2^p - 1$, and we omit the details.

Step 9 $K/H \not\cong {}^2F_4(q)$, where $q = 2^{2m+1} > 2$.

The odd order components of ${}^2F_4(q)$ are $q^2 + \sqrt{2q^3} + q + \sqrt{2q} + 1$ and $q^2 - \sqrt{2q^3} + q - \sqrt{2q} + 1$. Therefore, we have $q^2 + \sqrt{2q^3} + q + \sqrt{2q} + 1 = 2^p - 1$ or $q^2 - \sqrt{2q^3} + q - \sqrt{2q} + 1 = 2^p - 1$. However, if 2^{2m+1} is substituted in these equations we obtain $2^{m+1}(2^{3m+1} \pm 2^{2m+1} \pm 2^m \pm 1) = 2(2^{p-1} - 1)$, which is a contradiction.

Step 10 $K/H \not\cong F_4(q)$, where $q = 2^m$.

The odd order components of $F_4(q)$ are $q^4 + 1$ and $q^4 - q^2 + 1$. It is easy to see that in both cases, 2^{2m} divides $2(2^{p-1} - 1)$, a contradiction.

Step 11 $K/H \not\cong {}^2G_2(q)$, where $q = 3^{2m+1} > 3$.

The odd order components of ${}^2G_2(q)$ are $q + \sqrt{3q} + 1$ and $q - \sqrt{3q} + 1$. If $q - \sqrt{3q} + 1 = 2^p - 1$, then $q^3 > 2^{3p}$, while Lemma 8 shows that $q^3 < 2^{3p}$, which is a contradiction. If $q + \sqrt{3q} + 1 = 2^p - 1$, then

$$2^p - 2 = 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1) = 3^{m+1}(3^m + 1). \quad (1)$$

First of all, we recall that $(2^{\frac{p-1}{2}} - 1, 2^{\frac{p-1}{2}} + 1) = 1$. Now we consider two cases separately:

(i) If 3^{m+1} divides $2^{\frac{p-1}{2}} - 1$, then

$$3^m + 1 < 3^{m+1} \leq 2^{\frac{p-1}{2}} - 1 < 2^{\frac{p-1}{2}} + 1.$$

Hence, we obtain

$$3^{m+1}(3^m + 1) < 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1),$$

a contradiction.

(ii) If 3^{m+1} divides $2^{\frac{p-1}{2}} + 1$, then $2^{\frac{p-1}{2}} + 1 = k \cdot 3^{m+1}$ where k is a natural number. Now, from Eq. (1), it follows that

$$2k(2^{\frac{p-1}{2}} - 1) = 3^m + 1,$$

and consequently $3^m \geq 2^{\frac{p+1}{2}} - 1$. Therefore we have

$$2^{\frac{p+1}{2}} - 1 \leq 3^m < 3^{m+1} \leq 2^{\frac{p-1}{2}} + 1,$$

a contradiction.

Step 12 $K/H \not\cong G_2(q)$, where $q = 3^m$.

Recall that the odd order components of $G_2(q)$ are $q^2 - q + 1$ and $q^2 + q + 1$. If $q^2 - q + 1 = 2^p - 1$ then $q^6 > 2^{3p}$, while one can follow from Lemma 8 that $q^6 < 2^{3p}$, which is a contradiction. If $q^2 + q + 1 = 2^p - 1$, then $q(q + 1) \equiv 2 \pmod{4}$, which forces m is even. But then, it is obvious that $2^p - 2 = q(q + 1) \equiv 2 \pmod{8}$, a contradiction.

Step 13 $K/H \not\cong {}^2D_r(3)$, where $r = 2^m + 1$ is a prime number and $m \geq 1$.

Recall that

$$|{}^2D_r(3)| = \frac{1}{(4, 3^r + 1)} 3^{r(r-1)} (3^r + 1) \prod_{i=1}^{r-1} (3^{2i} - 1),$$

and the odd order components of ${}^2D_r(3)$ are

$$\frac{3^{r-1} + 1}{2} \quad \text{and} \quad \frac{3^r + 1}{4}.$$

In the case when $(3^{r-1} + 1)/2 = 2^p - 1$, adding 1 to both sides of this equality, we obtain $3(3^{r-2} + 1) = 2^{p+1}$, which is a contradiction. If $(3^r + 1)/4 = 2^p - 1$, then $r \geq 5$ because $p \geq 5$. Moreover, on the one hand, from last equation we obtain $3^r = 2^{p+2} - 5 > 2^{p+1}$, which implies that

$$3^{r(r-1)} > 2^{(p+1)(r-1)} > 2^{4(p+1)}.$$

On the other hand, it follows from Lemma 8 that

$$3^{r(r-1)} = |K/H|_3 \leq |G|_3 < 2^{3p},$$

which is a contradiction.

Step 14 $K/H \not\cong B_n(q)$, where $n = 2^m \geq 4$ and $q = r^f$ is an odd prime power.

Note that

$$|B_n(q)| = \frac{1}{(2, q-1)} q^{n^2} \prod_{i=1}^n (q^{2^i} - 1),$$

and the only odd order component of $B_n(q)$ is $(q^n+1)/2$. If $(q^n+1)/2 = 2^p-1$, then $q^n = 2^{p+1}-3 > 2^p$ and clearly q is not divisible by 2 and 3. Since $p \geq 5$ and $n \geq 4$, it is easy to see that

$$q^{n^2} > q^{3n} > 2^{3p} > 2^{2p}.$$

On the other hand, by Lemma 8, we obtain

$$q^{n^2} = |K/H|_r \leq |G|_r < 2^{2p},$$

which is a contradiction.

Step 15 $K/H \not\cong B_r(3)$.

The only odd order component of $B_r(3)$ is $(3^r-1)/2$. If $(3^r-1)/2 = 2^p-1$, then $2^{p+1}-3^r = 1$. However, this equation has no solution by Lemma 7, which is impossible.

Step 16 $K/H \not\cong {}^3D_4(q)$.

We recall that $q^4 - q^2 + 1$ is the only odd order component of ${}^3D_4(q)$, and so $q^4 - q^2 + 1 = 2^p - 1$. But then, $q^2(q^2 - 1) = 2(2^{p-1} - 1)$, which shows that $2(2^{p-1} - 1)$ is divisible by 4, a contradiction.

Step 17 $K/H \not\cong G_2(q)$, where $2 < q \equiv \pm 1 \pmod{3}$.

In this case, the odd order components of $G_2(q)$ are $q^2 + q + 1$ and $q^2 - q + 1$. Let $q = r^f$. If $q^2 + q + 1 = 2^p - 1$, then $q(q+1) = 2(2^{p-1} - 1)$, which shows that $q > 2$ is not a power of 2. Moreover, since $q - 1 \geq 2$, we obtain

$$q^3 - 1 = (q-1)(q^2 + q + 1) \geq 2(2^p - 1),$$

and so $q^3 \geq 2^{p+1} - 1 > 2^p$, which yields that $q^6 > 2^{2p}$. However, since $|G_2(q)| = q^6(q^2 - 1)(q^6 - 1)$, from Lemma 8, we conclude that

$$q^6 = |K/H|_r \leq |G|_r < 2^{2p},$$

which is a contradiction.

The case when $q^2 - q + 1 = 2^p - 1$ is similar and left to the reader.

Step 18 $K/H \not\cong {}^2D_n(3)$, where $n = 2^m + 1$ which is not a prime and $m \geq 2$.

The odd order component of ${}^2D_n(3)$ is $(3^{n-1}+1)/2$. If $(3^{n-1}+1)/2 = 2^p-1$, then $2^{p+1} = 3(3^{n-2}-1)$, a contradiction.

Step 19 $K/H \not\cong {}^2D_r(3)$, where $r \geq 5$ is a prime and $r \neq 2^m + 1$.

We recall that

$$|^2D_r(3)| = \frac{1}{(4, 3^r+1)} 3^{r(r-1)} (3^r+1) \prod_{i=1}^{r-1} (3^{2^i} - 1).$$

Moreover, the only odd order component of ${}^2D_r(3)$ is $(3^r+1)/4$. Let $(3^r+1)/4 = 2^p-1$. An easy computation shows that $3^r = 2^{p+2} - 5 > 2^{p+1}$. Moreover, we note that $r-1 \geq 4$, and so

$$3^{r(r-1)} \geq 3^{4r} > 2^{4(p+1)}.$$

On the other hand, by Lemma 8, we obtain

$$3^{r(r-1)} = |K/H|_3 \leq |G|_3 < 2^{3p},$$

which is a contradiction.

Step 20 $K/H \not\cong {}^2D_n(2)$, where $n = 2^m + 1$, $m \geq 2$.

The only odd order component of ${}^2D_n(2)$ is $2^{n-1} - 1$. If $2^{n-1} - 1 = 2^p - 1$, then $n - 1 = p$ and $2^m = p$, an impossible.

Step 21 $K/H \not\cong {}^2D_n(q)$, where $n = 2^m \geq 4$ and $q = r^f$.

Recall that

$$|{}^2D_n(q)| = \frac{1}{(4, q^n + 1)} q^{n(n-1)} (q^n + 1) \prod_{i=1}^{n-1} (q^{2^i} - 1),$$

and the only odd order component of ${}^2D_n(q)$ is $(q^n + 1)/(2, q + 1)$. Then $(q^n + 1)/(2, q + 1) = 2^p - 1$. Assume first that $(2, q + 1) = 1$. In this case, we obtain $q^n = 2(2^{p-1} - 1)$, a contradiction. Assume next that $(2, q + 1) = 2$. Again, using simple calculations we obtain $q^n = 2^{p+1} - 3 > 2^p$ and so q cannot be a power of 2. Thus, since $n - 1 \geq 3$, $q^{n(n-1)} \geq q^{3n} > 2^{3p}$. However, Lemma 8 shows that

$$q^{n(n-1)} = |K/H|_r \leq |G|_r < 2^{3p},$$

which is a contradiction.

Step 22 $K/H \not\cong D_{r+1}(q)$, where $q = 2, 3$.

The only odd order component of $D_{r+1}(q)$ is $(q^r - 1)/(2, q - 1)$, and so $(q^r - 1)/(2, q - 1) = 2^p - 1$. If $(2, q - 1) = 1$, then $r = p$ and $q = 2$, and we have

$$|K/H| = |D_{p+1}(2)| = \frac{1}{(4, 2^{p+1} - 1)} 2^{p(p+1)} (2^{p+1} - 1) \prod_{i=1}^p (2^{2^i} - 1),$$

this shows that $|K/H|_2$ does not divide $|G|_2$, which is a contradiction. In the case when $(2, q - 1) = 2$, we have the equation $2^{p+1} - 3^r = 1$, which has no solution for $p \geq 5$, by Lemma 7. This is the final contradiction.

Step 23 $K/H \not\cong D_r(q)$, where $q = 2, 3, 5$ and $r \geq 5$.

We recall that the only odd order component of $D_r(q)$ is $(q^r - 1)/(q - 1)$. We distinguish three cases separately.

(i) $q = 2$. In this case, we have $2^r - 1 = 2^p - 1$, and so $r = p$ and

$$|K/H| = |D_p(2)| = 2^{p(p-1)} (2^p - 1) \prod_{i=1}^{p-1} (2^{2^i} - 1).$$

Note that $|\text{Out}(D_p(2))| = 2$ and $D_p(2) \leq G/H \leq \text{Aut}(D_p(2))$. Now, considering the order of groups, we get $|H| = 2^\alpha (2^p + 1)$ where $p - 1 \leq \alpha \leq p$. Let $r \in \text{ppd}(2^{2^p} - 1)$ and $Q \in \text{Syl}_r(H)$. Clearly $r \in \pi(2^p + 1)$, Q is a normal π_1 -subgroup of G and $|Q|$ divides $2^p + 1$. Now, from Lemma 4, it follows that $|Q| - 1$ is divisible by $m_2(G) = 2^p - 1$, and so $|Q| - 1 \geq 2^p - 1$ or equivalently $|Q| \geq 2^p$. This forces $|Q| = 2^p + 1$. But then $m_2(G) = 2^p - 1$ does not divide the value $|Q| - 1 = 2^p$, which is a contradiction.

(ii) $q = 3$. In this case, from the equality $(3^r - 1)/2 = 2^p - 1$, we deduce that $2^{p+1} - 3^r = 1$. However, this equation has no solution when $p \geq 5$ by Lemma 7, a contradiction.

(iii) $q = 5$. Here $(5^r - 1)/4 = 2^p - 1$, and so $5^r = 2^{p+2} - 3 > 2^{p+1}$. As before, since $r - 1 \geq 4$, we obtain $5^{r(r-1)} > 5^{4r} > 2^{4(p+1)}$. On the other hand, we have

$$5^{r(r-1)} = |K/H|_5 \leq |G|_5 < 2^{2p},$$

by Lemma 8, which is a contradiction.

Step 24 $K/H \not\cong C_r(3)$.

The only odd order component of $C_r(3)$ is $(3^r - 1)/2$. Thus, if $(3^r - 1)/2 = 2^p - 1$, then $2^{p+1} - 3^r = 1$. However, this equation has no solution by Lemma 7, an impossible.

Step 25 $K/H \not\cong C_n(q)$, where $n = 2^m \geq 2$.

Note that

$$|C_n(q)| = \frac{1}{(2, q-1)} q^{n^2} \prod_{i=1}^n (q^{2^i} - 1),$$

and the only odd order component of $C_n(q)$ is $(q^n + 1)/(2, q-1)$. Thus $(q^n + 1)/(2, q-1) = 2^p - 1$. If $(2, q-1) = 1$, then $q^n = 2(2^{p-1} - 1)$, which yields that $q = p = 2$ and $n = 1$, a contradiction. If $(2, q-1) = 2$, then $q^n = 2^{p+1} - 3 > 2^p$, which implies that q is not a power of 2 and 3. Let $q = r^f$. When $n \geq 4$, it is easy to see that

$$q^{n^2} > q^{3n} > 2^{3p} > 2^{2(p+1)}.$$

But, from Lemma 8, we obtain

$$q^{n^2} = |K/H|_r \leq |G|_r < 2^{2p},$$

a contradiction. Assume now that $n = 2$. In this case, we have $q^2 = 2^{p+1} - 3$, or equivalently

$$(q-1)(q+1) = 2^2(2^{p-1} - 1).$$

However, the left hand side is divisible by 8, while the right hand side is divisible by 4, a contradiction.

Step 26 $K/H \not\cong A_1(q)$, where $q = 2^m > 2$.

The odd order components of $A_1(q)$ are $q+1$ and $q-1$. If $q+1 = 2^p - 1$, then $q = 2(2^{p-1} - 1)$, a contradiction. If $q-1 = 2^p - 1$, then $q = 2^p$. Moreover, since $A_1(q) \leq G/H \leq \text{Aut}(A_1(q))$, it is easy to see that the order of H is divisible by $(2^2 - 1)(2^4 - 1) \cdots (2^{2(p-1)} - 1)$. Let $r \in \text{ppd}(2^{2(p-1)} - 1)$ and $Q \in \text{Syl}_r(H)$. Clearly Q is a π_1 -normal subgroup of G and $|Q|$ divides $2^{p-1} + 1$. On the other hand, from Lemma 4, $|Q| - 1$ is divisible by $2^p - 1$ which implies that $|Q| \geq 2^p$. This is a contradiction.

Step 27 $K/H \not\cong A_1(q)$, where $3 \leq q \equiv \pm 1 \pmod{4}$ and $q = r^f$.

Assume first that $3 \leq q \equiv 1 \pmod{4}$. In this case, the odd order components of $A_1(q)$ are $(q+1)/2$ and q . If $(q+1)/2 = 2^p - 1$, then $r^f = q = 2^{p+1} - 3$. First of all, we claim that f is an odd number. Otherwise, we have

$$(r^{\frac{f}{2}} - 1)(r^{\frac{f}{2}} + 1) = 2^2(2^{p-1} - 1).$$

But then, the left hand side is divisible by 8, while the right hand side is divisible by 4, which is a contradiction. Furthermore, by easy computations we observe that

$$|A_1(q)| = \frac{1}{2}q(q^2 - 1) = 2^2(2^{p+1} - 3)(2^{p-1} - 1)(2^p - 1).$$

On the other hand, since $|G/K| \cdot |H| = |G|/|A_1(q)|$, we deduce that

$$|G/K|_2 \cdot |H|_2 = \frac{|G|_2}{|A_1(q)|_2} = 2^{p^2-2}.$$

But since $|G/K|$ divides $|\text{Out}(A_1(q))| = 2f$ and f is odd, $|G/K|_2$ is at most 2. Hence, if $S_2 \in \text{Syl}_2(H)$, then $|S_2| = 2^{p^2-2}$ or $|S_2| = 2^{p^2-3}$. We notice that S_2 is a normal subgroup of G , because H is nilpotent. Now, it follows from Lemma 4 that $2^p - 1$ divides $2^{p^2-2} - 1$ or $2^{p^2-3} - 1$, which is a contradiction. If $q = 2^p - 1$, we get a contradiction by Lemma 7.

Assume next that $3 \leq q \equiv -1 \pmod{4}$. In this case, the odd order components of $A_1(q)$ are $(q-1)/2$ and q . If $(q-1)/2 = 2^p - 1$, then $2^{p+1} - r^f = 1$. Noting Lemma 7, we deduce that $f = 1$, and hence $r = 2^{p+1} - 1$ is a Mersenne prime, which is a contradiction because $p+1$ is not a prime. The case when $q = 2^p - 1$ is similar to the previous paragraph.

Step 28 $K/H \not\cong A_r(q)$, where $(q-1) \mid (r+1)$.

Recall that

$$|K/H| = |A_r(q)| = \frac{1}{(r+1, q-1)} q^{r(r+1)/2} \prod_{i=2}^{r+1} (q^i - 1),$$

and the only odd order component of $A_r(q)$ is $(q^r - 1)/(q - 1)$, and so $(q^r - 1)/(q - 1) = 2^p - 1$. As a simple observation we see that $q^r - 1 \geq (q^r - 1)/(q - 1) = 2^p - 1$ and so $q^r \geq 2^p$. Let $q = t^f$, where t is a prime number and f is a natural number.

- (i) Suppose first that $r \geq 7$. Then $q^{\frac{r(r+1)}{2}} > q^{3(r+1)} \geq 2^3 q^{3r} \geq 2^{3(p+1)}$. Now, if t is an odd prime, then by Lemma 8 we obtain

$$q^{r(r+1)/2} = |K/H|_t \leq |G|_t < 2^{3p},$$

which is a contradiction. Therefore, we may assume that $t = 2$. In this case, we have $(2^{fr} - 1)/(2^f - 1) = 2^p - 1$, from which one can deduce that $f = 1$ and $r = p$. Thus

$$|G/K| \cdot |H| = \frac{2^{p^2} \prod_{i=1}^p (2^{2i} - 1)}{2^{\frac{p(p+1)}{2}} \prod_{i=2}^{p+1} (2^i - 1)}.$$

Since $|G/K|$ divides $|\text{Out}(K/H)| = |\text{Out}(A_p(2))| = 2$, we conclude that $|H|$ is divisible by $2^p + 1$. Let $s \in \text{ppd}(2^{2p} - 1) \subseteq \pi(2^p + 1)$ and $Q \in \text{Syl}_s(H)$. Clearly $|Q| \mid 2^p + 1$. Since H is a normal π_1 -subgroup of G which is nilpotent, Q is also a normal π_1 -subgroup of G . Now, by Lemma 4, $m_2(G) = 2^p - 1$ divides $|Q| - 1$, and so $|Q| \geq 2^p$. But, this forces $|Q| = 2^p + 1$. However, this contradicts the fact that $2^p - 1 \mid 2^p$.

- (ii) Suppose next that $r = 5$. If q is even, then from $(q^5 - 1)/(q - 1) = 2^p - 1$, we obtain $q(q^3 + q^2 + q + 1) = 2(2^{p-1} - 1)$, which implies that $q = 2$ and $r = p = 5$. Therefore, by easy calculations we see that

$$|G/K| \cdot |H| = \frac{2^{10} \prod_{i=1}^5 (2^i + 1)}{2^6 - 1},$$

which is not a natural number, a contradiction. If q is odd, then we get

$$q(q+1)(q^2+1) = q^4 + q^3 + q^2 + q = 2^p - 2,$$

however $q(q+1)(q^2+1) \equiv 0 \pmod{4}$, while $2^p - 2 \equiv 2 \pmod{4}$, a contradiction.

- (iii) Finally suppose that $r = 3$. Then $q(q+1) = 2(2^{p-1} - 1)$. First of all, we note that q is not even, otherwise $p = 3$, an impossible. In addition, we have

$$q(q+1) = 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1). \quad (2)$$

Now we consider two cases separately:

- (a) If q divides $2^{\frac{p-1}{2}} - 1$, then

$$q \leq 2^{\frac{p-1}{2}} - 1, \quad q+1 < 2^{\frac{p-1}{2}} + 1.$$

Hence, we obtain

$$q(q+1) < 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1),$$

a contradiction.

(b) If q divides $2^{\frac{p-1}{2}} + 1$, then $2^{\frac{p-1}{2}} + 1 = kq$ for some natural number k . Now from Eq. (2), it follows that

$$2k(2^{\frac{p-1}{2}} - 1) = q + 1.$$

If $k = 1$, then $p = q = 5$. Hence $13 \in \pi(K/H) = \pi(A_3(5))$ but $13 \notin \pi(G) = \pi(C_5(2))$, a contradiction. Thus $k \geq 2$ and we obtain

$$2(2^{\frac{p+1}{2}} - 2) - 1 \leq q < q + 1 \leq kq = 2^{\frac{p-1}{2}} + 1,$$

a contradiction.

Step 29 $K/H \not\cong A_{r-1}(q)$, where $(r, q) \neq (3, 2), (3, 4)$.

Again, we recall that

$$|K/H| = |A_{r-1}(q)| = \frac{1}{(r, q-1)} q^{r(r-1)/2} \prod_{i=2}^r (q^i - 1),$$

and the only odd order component of $A_{r-1}(q)$ is $(q^r - 1)/(q - 1)(r, q - 1)$. Hence, we must have

$$(q^r - 1)/(q - 1)(r, q - 1) = 2^p - 1,$$

which implies that

$$q^r - 1 \geq (q^r - 1)/(q - 1)(r, q - 1) = 2^p - 1,$$

or equivalently $q^r \geq 2^p$. Let $q = t^f$, where t is a prime and f is a natural number. In what follows, we consider several cases separately.

(i) $r \geq 7$. In this case, we obtain

$$q^{r(r-1)/2} \geq q^{3r} \geq 2^{3p},$$

and Lemma 8 implies that $t = 2$. Now, Lemma 1 shows that $q = 2$ and $r = p$, and hence we obtain

$$|G/K| \cdot |H| = \frac{2^{p^2} \prod_{i=1}^p (2^{2i} - 1)}{2^{\binom{p}{2}} \prod_{i=2}^p (2^i - 1)} = 2^{\frac{p(p+1)}{2}} \prod_{i=1}^p (2^i + 1).$$

On the other hand, $|G/K|$ divides $|\text{Out}(K/H)| = 2$. From this we deduce that $|H|$ is divisible by $2^p + 1$. Let $s \in \text{ppd}(2^{2p} - 1) \subseteq \pi(2^p + 1)$ and $Q \in \text{Syl}_s(H)$. Evidently Q is a normal subgroup of G and $|Q|$ divides $2^p + 1$. Now, it follows from Lemma 4 that $m_2(G) = 2^p - 1 \mid |Q| - 1$, which is impossible.

(ii) $r = 5$. Assume first that $(5, q - 1) = 1$. In this case, we have

$$\frac{q^5 - 1}{q - 1} = q^4 + q^3 + q^2 + q + 1 = 2^p - 1,$$

or equivalently

$$q(q + 1)(q^2 + 1) = 2(2^{p-1} - 1). \quad (3)$$

If q is even, then we conclude that $q = 2$ and $r = p = 5$, and the proof is quite similar as (i). If q is odd, then the left-hand side of Eq. (3) is congruent to 0 (mod 4), while the right-hand side of Eq. (3) is congruent to 2 (mod 4), a contradiction.

Assume next that $(5, q - 1) = 5$. In this case, we have

$$q^4 + q^3 + q^2 + q + 1 = 5(2^p - 1),$$

or equivalently

$$(q-1)(q^3 + 2q^2 + 3q + 4) = 10(2^{p-1} - 1).$$

In the case when q is even, it follows that $q = 2$ and so $13 = 5(2^{p-1} - 1)$, a contradiction. Moreover, if q is odd, then from the equality $q(q+1)(q^2+1) = 5 \cdot 2^p - 6$ it is easily seen that the left-hand side of this equation is congruent to 0 (mod 4), while the right-hand side is congruent to 2 (mod 4), a contradiction.

(iii) $r = 3$. In this case, we have $(q^3 - 1)/(q-1)(3, q-1) = 2^p - 1$. First of all, if q is even, then we obtain $p = 3$, which is not the case. Thus, we can assume that q is odd.

If $(3, q-1) = 1$, then

$$q(q+1) = 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1). \quad (4)$$

If q divides $2^{\frac{p-1}{2}} - 1$, then

$$q \leq 2^{\frac{p-1}{2}} - 1, \quad q+1 < 2^{\frac{p-1}{2}} + 1.$$

Hence, we obtain

$$q(q+1) < 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1),$$

a contradiction. If q divides $2^{\frac{p-1}{2}} + 1$, then $2^{\frac{p-1}{2}} + 1 = kq$. Now, from Eq. (4), it follows that

$$2k(2^{\frac{p-1}{2}} - 1) = q + 1.$$

When $k = 1$, we conclude that $p = 5$ and $q = 5$. But then, we have $|K/H| = |A_2(5)| = 2^5 \cdot 3 \cdot 5^3 \cdot 31$, while $|G| = |C_5(2)| = 2^{25} \cdot 3^6 \cdot 5^2 \cdot 7 \cdot 11 \cdot 17 \cdot 31$, this is a contradiction because $|K/H|_5 > |G|_5$. If $k \geq 2$, then $q \geq 2(2^{\frac{p-1}{2}} - 2) - 1$. Therefore, we have

$$2(2^{\frac{p+1}{2}} - 2) - 1 \leq q < q+1 \leq 2^{\frac{p-1}{2}} + 1,$$

a contradiction.

If $(3, q-1) = 3$, then $q(q+1) = 2^2(3 \cdot 2^{p-2} - 1)$, which implies that $4 \parallel q+1$ and so $2 \parallel q-1$. Moreover, under these conditions, one can easily deduce that f is odd, otherwise $8 \parallel q-1 = t^f - 1 = (t^{\frac{f}{2}} - 1)(t^{\frac{f}{2}} + 1)$, which is a contradiction. Thus, we have $|A_2(q)|_2 = 2^4$, while

$$|G/K|_2 \cdot |H|_2 = \frac{|G|_2}{|A_2(q)|_2} = 2^{p^2-4}.$$

Since $|G/K|$ divides $2f(3, q-1)$ and f is odd, $|G/K|_2 \leq 2$. Therefore a Sylow 2-subgroup of H has order either 2^{p^2-4} or 2^{p^2-5} . Applying Lemma 4 we deduce that $2^p - 1 \mid 2^{p^2-4} - 1$ or $2^p - 1 \mid 2^{p^2-5} - 1$. Now, one can easily check that the second divisibility is possible only for $p = 5$. But then, we get $q(q+1) = 2^2 \cdot 23$, which is a contradiction.

Step 30 $K/H \not\cong {}^2A_r(q)$, where $(q+1) \mid (r+1)$ and $(r, q) \neq (3, 3), (5, 2)$.

In this case, we have

$$|K/H| = |{}^2A_r(q)| = \frac{1}{(r+1, q+1)} q^{r(r+1)/2} \prod_{i=2}^{r+1} (q^i - (-1)^i),$$

and the odd order component of ${}^2A_r(q)$ is $(q^r + 1)/(q + 1)$, and so $(q^r + 1)/(q + 1) = 2^p - 1$. An argument similar to that in the previous cases shows that $q^r - 1 > (q^r + 1)/(q + 1) = 2^p - 1$, and so $q^r > 2^p$. Let $q = t^f$. We now consider three cases separately.

- (i) $r \geq 7$. Then $q^{\frac{r(r+1)}{2}} > q^{3(r+1)} \geq 2^3 q^{3r} > 2^{3(p+1)}$, which forces by Lemma 8 that $t = 2$. Thus $(2^{fr} + 1)/(2^f + 1) = 2^p - 1$, and, consequently, $f = 1$, $r = 3$ and $p = 2$, which is a contradiction.
- (ii) If $r = 5$, then $(q^5 + 1)/(q + 1) = 2^p - 1$. Arguing as in the case (i), we conclude that $t = 2$ and $f = 1$, whence $12 = 2^p$, a contradiction.
- (iii) If $r = 3$, then $(q^3 + 1)/(q + 1) = 2^p - 1$. It follows that $q(q - 1) = 2(2^{p-1} - 1)$, and so $q = p = 2$, which is impossible.

Step 31 $K/H \not\cong {}^2A_{r-1}(q)$.

In this case, we have

$$|K/H| = |{}^2A_{r-1}(q)| = \frac{1}{(r, q+1)} q^{r(r-1)/2} \prod_{i=2}^r (q^i - (-1)^i),$$

and the odd order component of ${}^2A_{r-1}(q)$ is $(q^r + 1)/(q + 1)(r, q + 1)$. Thus

$$\frac{q^r + 1}{(q + 1)(r, q + 1)} = 2^p - 1,$$

As before, we deduce that $q^r \geq 2^p$. Let $q = t^f$. We now consider three cases separately.

- (i) $r \geq 7$. It follows that $q^{r(r-1)/2} \geq q^{3r} > 2^{3p}$, which implies that $t = 2$ by Lemma 8. Now, we obtain $\frac{2^{fr} + 1}{(2^f + 1)(r, 2^f + 1)} = 2^p - 1$, which contradicts Lemma 1 because $2^p - 1$ is the largest prime in $\pi(G)$.
- (ii) $r = 5$. In this case we have $q^5 + 1 = (q + 1)(2^p - 1)(5, q + 1)$.
Assume first that q is even, that is $q = 2^f$. If $(5, q + 1) = 1$, then we obtain $2^{5f} = 2^{fp} + 2^p - 2^f - 2$, which is impossible. If $(5, q + 1) = 5$, then $2^{5f} = 5(2^{fp} + 2^p - 2^f) - 6$, which is again a contradiction.
Assume next that q is odd. Noting that $q(q - 1)(q^2 + 1) = (2^p - 1)(5, q + 1) - 1$, it is easily seen that the left hand side is congruent to 0 (mod 4), while the right hand side is congruent to 2 (mod 4), a contradiction.
- (iii) $r = 3$. In this case, we have $(q^3 + 1)/(q + 1)(3, q + 1) = 2^p - 1$. If $(3, q + 1) = 1$, then we obtain

$$q(q - 1) = 2^p - 2 = 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1).$$

If q divides 2, then $p = 2$, a contradiction. If q divides $2^{\frac{p-1}{2}} - 1$ or $2^{\frac{p-1}{2}} + 1$, then

$$q(q - 1) < 2^p - 2 = 2(2^{\frac{p-1}{2}} - 1)(2^{\frac{p-1}{2}} + 1),$$

a contradiction. Therefore we may assume that $(3, q + 1) = 3$. If q is even, then we conclude that $q = 4$, which is a contradiction. We now suppose that q is odd. Since $q(q - 1) = 2^2(3 \cdot 2^{p-2} - 1)$, it follows that $4|q - 1$, and so $2||q + 1$. Moreover, under these hypotheses, one can easily deduce that f is odd, otherwise $8|q - 1 = t^f - 1 = (t^{\frac{f}{2}} - 1)(t^{\frac{f}{2}} + 1)$, which is a contradiction. On the other hand, $|G/K|$ divides $f(3, q + 1)$ and since f is odd, $|G/K|_2 = 1$. Therefore a Sylow 2-subgroup of H has order 2^{p^2-4} . Again, using Lemma 4, we see that $2^p - 1 | 2^{p^2-4} - 1$, which implies that $p = 2$. This is a contradiction.

Step 32 $K/H \not\cong C_r(2)$.

The odd order component of $C_r(2)$ is $2^r - 1$. Thus $2^r - 1 = 2^p - 1$. It follows that $r = p$, $G/K = 1$ and $H = 1$, which means $G \cong C$. This completes the proof of the theorem. $\square$

4 Appendix

In a series of articles, it was shown that many finite simple groups are OD-characterizable or 2-fold OD-characterizable. Table 1 lists finite simple groups which are currently known to be k -fold OD-characterizable for $k \in \{1, 2\}$. Until recently, no examples of simple groups P with $h_{\text{OD}}(P) \geq 3$ were known. Therefore, we posed the following question:

Problem 1 *Is there a non-abelian simple group P with $h_{\text{OD}}(P) \geq 3$?*

Table 1. Some non-abelian simple groups S with $h_{\text{OD}}(S) = 1$ or 2.

S	Conditions on S	$h_{\text{OD}}(S)$	Refs.
$\mathbb{A}_n$	$n = p, p+1, p+2$ (p a prime)	1	[18], [21]
	$5 \leq n \leq 100, n \neq 10$	1	[9], [11], [16], [19], [30]
	$n = 106, 112$	1	[27]
	$n = 10$	2	[20]
$L_2(q)$	$q \neq 2, 3$	1	[18], [21], [36]
$L_3(q)$	$ \pi(\frac{q^2+q+1}{d}) = 1, d = (3, q-1)$	1	[21]
$U_3(q)$	$ \pi(\frac{q^2-q+1}{d}) = 1, d = (3, q+1), q > 5$	1	[21]
$L_3(9)$		1	[33]
$U_3(5)$		1	[34]
$L_4(q)$	$q \leq 17$	1	[1, 3]
$U_4(7)$		1	[3]
$L_n(2)$	$n = p$ or $p+1$, for which $2^p - 1$ is a prime	1	[3]
$L_n(2)$	$n = 9, 10, 11$	1	[10], [17]
$U_6(2)$		1	[35]
$R(q)$	$ \pi(q \pm \sqrt{3q} + 1) = 1, q = 3^{2m+1}, m \geq 1$	1	[21]
$\text{Sz}(q)$	$q = 2^{2n+1} \geq 8$	1	[18], [21]
$B_m(q), C_m(q)$	$m = 2^f \geq 4, \pi((q^m + 1)/2) = 1,$	2	[2]
$B_2(q) \cong C_2(q)$	$ \pi((q^2 + 1)/2) = 1, q \neq 3$	1	[2]
$B_m(q) \cong C_m(q)$	$m = 2^f \geq 2, 2 q, \pi(q^m + 1) = 1, (m, q) \neq (2, 2)$	1	[2]
$B_p(3), C_p(3)$	$ \pi((3^p - 1)/2) = 1, p$ is an odd prime	2	[2], [21]
$B_3(5), C_3(5)$		2	[2]
$C_3(4)$		1	[14]
S	A sporadic simple group	1	[21]
S	A simple group with $ \pi(S) = 4, S \neq \mathbb{A}_{10}$	1	[32]
S	A simple group with $ S \leq 10^8, S \neq \mathbb{A}_{10}, U_4(2)$	1	[29]
S	A simple $C_{2,2}$ -group	1	[18]

Although we have not found a simple group which is k -fold OD-characterizable for $k \geq 3$, but among non-simple groups, there are many groups which are k -fold OD-characterizable for $k \geq 3$. As

In Table 3, q is a power of a prime number.

an easy example, if P is a p -group of order p^n , then $h_{\text{OD}}(P) = \nu(p^n)$, where $\nu(m)$ signifies the number of non-isomorphic groups of order m . Table 2 lists finite non-solvable groups which are currently known to be OD-characterizable or k -fold OD-characterizable with $k \geq 2$.

Table 2. Some non-solvable groups G with known $h_{\text{OD}}(G)$.

G	Conditions on G	$h_{\text{OD}}(G)$	Refs.
$\text{Aut}(M)$	M is a sporadic group $\neq J_2, M^cL$	1	[18]
$\mathbb{S}_n$	$n = p, p+1$ ($p \geq 5$ is a prime)	1	[18]
$U_3(5) : 2$		1	[34]
$U_6(2) : 2$		1	[35]
M	$M \in \mathcal{C}_1$	2	[20]
M	$M \in \mathcal{C}_2$	8	[20]
M	$M \in \mathcal{C}_3$	3	[9, 11, 16, 19, 27]
M	$M \in \mathcal{C}_4$	2	[20]
M	$M \in \mathcal{C}_5$	3	[20]
M	$M \in \mathcal{C}_6$	6	[16]
M	$M \in \mathcal{C}_7$	1	[31]
M	$M \in \mathcal{C}_8$	9	[31]
M	$M \in \mathcal{C}_9$	3	[34]
M	$M \in \mathcal{C}_{10}$	6	[34]
M	$M \in \mathcal{C}_{11}$	3	[35]
M	$M \in \mathcal{C}_{12}$	5	[35]
M	$M \in \mathcal{C}_{13}$	1	[28]
M	$M \in \mathcal{C}_{14}$	1	[17]

$$\mathcal{C}_1 = \{\mathbb{A}_{10}, J_2 \times \mathbb{Z}_3\}$$

$$\mathcal{C}_2 = \{\mathbb{S}_{10}, \mathbb{Z}_2 \times \mathbb{A}_{10}, \mathbb{Z}_2 \cdot \mathbb{A}_{10}, \mathbb{Z}_6 \times J_2, \mathbb{S}_3 \times J_2, \mathbb{Z}_3 \times (\mathbb{Z}_2 \cdot J_2), (\mathbb{Z}_3 \times J_2) \cdot \mathbb{Z}_2, \mathbb{Z}_3 \times \text{Aut}(J_2)\}.$$

$$\mathcal{C}_3 = \{\mathbb{S}_n, \mathbb{Z}_2 \cdot \mathbb{A}_n, \mathbb{Z}_2 \times \mathbb{A}_n\}, \text{ where } 9 \leq n \leq 100 \text{ with } n \neq 10, p, p+1 \text{ (} p \text{ a prime)} \\ \text{or } n = 106, 112.$$

$$\mathcal{C}_4 = \{\text{Aut}(M^cL), \mathbb{Z}_2 \times M^cL\}.$$

$$\mathcal{C}_5 = \{\text{Aut}(J_2), \mathbb{Z}_2 \times J_2, \mathbb{Z}_2 \cdot J_2\}.$$

$$\mathcal{C}_6 = \{\text{Aut}(S_6(3)), \mathbb{Z}_2 \times S_6(3), \mathbb{Z}_2 \cdot S_6(3), \mathbb{Z}_2 \times O_7(3), \mathbb{Z}_2 \cdot O_7(3), \text{Aut}(O_7(3))\}.$$

$$\mathcal{C}_7 = \{L_2(49) : 2_1, L_2(49) : 2_2, L_2(49) : 2_3\}.$$

$$\mathcal{C}_8 = \{L \cdot 2^2, \mathbb{Z}_2 \times (L : 2_1), \mathbb{Z}_2 \times (L : 2_2), \mathbb{Z}_2 \times (L : 2_3), \mathbb{Z}_2 \cdot (L : 2_1), \\ \mathbb{Z}_2 \cdot (L : 2_2), \mathbb{Z}_2 \cdot (L : 2_3), \mathbb{Z}_4 \times L, (\mathbb{Z}_2 \times \mathbb{Z}_2) \times L\}, \text{ where } L = L_2(49).$$

$$\mathcal{C}_9 = \{U_3(5) : 3, \mathbb{Z}_3 \times U_3(5), \mathbb{Z}_3 \cdot U_3(5)\}$$

$$\mathcal{C}_{10} = \{L : \mathbb{S}_3, \mathbb{Z}_2 \cdot (L : 3), \mathbb{Z}_3 \times (L : 2), \mathbb{Z}_3 \cdot (L : 2), (\mathbb{Z}_2 \times L) : 2, (\mathbb{Z}_3 \cdot L) : 2\}, \\ \text{where } L = U_3(5).$$

$$\mathcal{C}_{11} = \{U_6(2) : 3, \mathbb{Z}_3 \times U_6(2), \mathbb{Z}_3 \cdot U_6(2)\}.$$

$$\mathcal{C}_{12} = \{L \cdot \mathbb{S}_3, \mathbb{Z}_3 \times (L : 2), \mathbb{Z}_3 \cdot (L : 2), (\mathbb{Z}_3 \times L) : 2, (\mathbb{Z}_3 \cdot L) : 2\}, \text{ where } L = U_6(2).$$

$$\mathcal{C}_{13} = \{\text{Aut}(O_{10}^+(2), \text{Aut}(O_{10}^-(2)),$$

$$\mathcal{C}_{14} = \{\text{Aut}(L_p(2)), \text{Aut}(L_{p+1}(2))\}, \text{ where } 2^p - 1 \text{ is a Mersenne prime.}$$

References

- [1] B. Akbari and A. R. Moghaddamfar, *Recognizing by order and degree pattern of some projective special linear groups*, International Journal of Algebra and Computation, 22(6)(2012), 22 pages.
- [2] M. Akbari and A. R. Moghaddamfar, *Simple groups which are 2-fold OD-characterizable*, Bull. Malays. Math. Sci. Soc., 35(1)(2012), 65-77.
- [3] M. Akbari, A. R. Moghaddamfar and S. Rahbariyan, *A characterization of some finite simple groups through their orders and degree patterns*, Algebra Colloq., 19(3)(2012), 473-482.
- [4] R. W. Carter, *Simple groups of Lie type*, Wiley, London (1972).
- [5] G. Y. Chen, *A new characterization of sporadic simple groups*, Algebra Colloq., 3(1)(1996), 49-58.
- [6] G. Y. Chen, *A new characterization of sporadic groups*, Algebra Colloq., 3(1)(1996) 49-58.
- [7] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, *Atlas of finite groups*, Clarendon Press, oxford, 1985.
- [8] P. Crescenzo, *A Diophantine equation which arises in the theory finite groups*, Advances in Math., 17 (1975), 25-29.
- [9] A. A. Hoseini and A. R. Moghaddamfar, *Recognizing alternating groups A_{p+3} for certain primes p by their orders and degree patterns*, Front. Math. China, 5(3)(2010), 541-553.
- [10] B. Khosravi, *Some characterizations of $L_9(2)$ related to its prime graph*, Publicationes Mathematicae Debrecen, 75(3-4)(2009), 375-385.
- [11] R. Kogani-Moghaddam and A. R. Moghaddamfar, *Groups with the same order and degree pattern*, Science China Mathematics, 55(4)(2012), 701-720.
- [12] A. S. Kondratév, *On prime graph components of finite simple groups*, Math. Sb., 180(6)(1989), 787-797.
- [13] A. S. Kondratév and V. D. Mazurov, *Recognition of alternating groups of prime degree from their element orders*, Siberian Mathematical Journal, 41(2)(2000), 294-302.
- [14] A. R. Moghaddamfar, *Recognizability of finite groups by order and degree pattern*, Proceedings of the International Conference on Algebra 2010, 422-433.
- [15] A. R. Moghaddamfar, *A comparison of the order components in Frobenius and 2-Frobenius groups with finite simple groups*, Taiwanese Journal of Mathematics, 13(1)(2009), 67-89.
- [16] A. R. Moghaddamfar and S. Rahbariyan, *More on the OD-characterizability of a finite group*, Algebra Colloquium, 18(4)(2011), 663-674.
- [17] A. R. Moghaddamfar and S. Rahbariyan, *OD-characterization of some linear groups over binary field and their automorphism groups*, Submitted for publication.
- [18] A. R. Moghaddamfar and A. R. Zokayi, *Recognizing finite groups through order and degree pattern*, Algebra Colloquium, 15(3)(2008), 449-456.
- [19] A. R. Moghaddamfar and A. R. Zokayi, *OD-characterization of alternating and symmetric groups of degrees 16 and 22*, Front. Math. China, 4(4)(2009), 669-680.

- [20] A. R. Moghaddamfar and A. R. Zokayi, *OD-characterization of certain finite groups having connected prime graphs*, Algebra Colloquium, 17(1)(2010), 121-130.
- [21] A. R. Moghaddamfar, A. R. Zokayi and M. R. Darafsheh, *A characterization of finite simple groups by the degrees of vertices of their prime graphs*, Algebra Colloquium, 12(3)(2005), 431-442.
- [22] H. Shi and G. Y. Chen, ${}^2D_{p+1}(2)$ ($5 \leq p \neq 2^m - 1$) can be characterized by its order components, Kumamoto J. Math., 18 (2005), 1-8.
- [23] M. Suzuki, *On the prime graph of a finite simple group—An application of the method of Feit-Thompson-Bender-Glauberman*, Groups and combinatorics—in memory of Michio Suzuki, Adv. Stud. Pure Math., 32 (Math. Soc. Japan, Tokyo, 2001), 41-207.
- [24] A. V. Vasilév and E. P. Vdovin, *An adjacency criterion in the prime graph of a finite simple group*, Algebra and Logic, 44(6)(2005), 381-406.
- [25] A. V. Vasilév and E. P. Vdovin, *Cocliques of maximal size in the prime graph of a finite simple group*, Algebra and Logic, 50(4)(2011), 291-322.
- [26] J. S. Williams, *Prime graph components of finite groups*, J. Algebra, 69(2)(1981), 487-513.
- [27] Y. X. Yan and G. Y. Chen, *OD-characterization of alternating and symmetric groups of degree 106 and 112*, Proceedings of the International Conference on Algebra 2010, 690-696.
- [28] Y. X. Yan, G. Y. Chen and L. L. Wang, *OD-characterization of the automorphism groups of $O_{10}^{\pm}(2)$* , Indian J. Pure Appl. Math., 43(3)(2012), 183-195.
- [29] L. C. Zhang and W. J. Shi, *OD-characterization of all simple groups whose orders are less than 10^8* , Front. Math. China, 3(3)(2008), 461-474.
- [30] L. C. Zhang, W. J. Shi, L. L. Wang and C. G. Shao, *OD-characterization of A_{16}* , Journal of Suzhou University (Natural Science Edition), 24(2)(2008), 7-10.
- [31] L. C. Zhang and W. J. Shi, *OD-characterization of almost simple groups related to $L_2(49)$* , Arch. Math. (Brno), 44(3)(2008), 191-199.
- [32] L. C. Zhang and W. J. Shi, *OD-characterization of simple K_4 -groups*, Algebra Colloq., 16(2)(2009), 275-282.
- [33] L. C. Zhang, W. J. Shi, C. G. Shao and L. L. Wang, *OD-characterization of the simple group $L_3(9)$* , Journal of Guangxi University (Natural Science Edition), 34(1)(2009), 120-122.
- [34] L. C. Zhang and W. J. Shi, *OD-characterization of almost simple groups related to $U_3(5)$* , Acta Mathematica Sinica (English Series), 26(1)(2010), 161-168.
- [35] L. C. Zhang and W. J. Shi, *OD-characterization of almost simple groups related to $U_6(2)$* , Acta Math. Sci. Ser. B, 31(2)(2011), 441-450.
- [36] L. C. Zhang and W. J. Shi, *OD-characterization of the projective special linear groups $L_2(q)$* , Algebra Colloq., 19(3)(2012), 509-524.
- [37] K. Zsigmondy, *Zur theorie der potenzreste*, Monatsh. Math. Phys., 3 (1892), 265-284.